Handout for Lecture 12

Good Estimators, Sample Mean Distribution, and Confidence Intervals

ECON 340: Economic Research Methods

Instructor: Div Bhagia

Good Estimators

Denote $\hat{\theta}$ as an estimator for the population parameter θ . Some desirable properties for an estimator

- Unbiasedness: $E(\hat{\theta}) = \theta$
- Efficiency: lower variance is better
- Consistency: as the sample size becomes infinitely large, $\hat{\theta} \rightarrow \theta$

Question 1: If some sample estimator $\hat{\theta}$ is an unbiased estimator for the true population parameter θ i.e. $E(\hat{\theta}) = \theta$. This implies that:

- $\Box \hat{\theta} = \theta$ in all samples.
- \Box If we take repeated samples, average of $\hat{\theta}$ is equal to θ

Question 2: We are choosing between two estimators $\hat{\theta}_1$ and $\hat{\theta}_2$, both of which are unbiased i.e. $E(\hat{\theta}_1) = \mu$ and $E(\hat{\theta}_2) = \mu$. But the variance of $\hat{\theta}_1$ is lower than that of $\hat{\theta}_2$ i.e. $Var(\hat{\theta}_1) < Var(\hat{\theta}_2)$. Which of the following is true?

- $\hfill\square$ We are indifferent between the two estimators.
- \Box We prefer $\hat{\theta}_1$ over $\hat{\theta}_2$.
- \Box We prefer $\hat{\theta}_2$ over $\hat{\theta}_1$.
- □ We need more information to reach any conclusion.

Sample Mean Distribution

Let $X_1, X_2, ..., X_n$ denote independent random draws (random sample) from a population with mean μ and variance σ^2 . Then the sample mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$ is a random variable with:

$$E(\bar{X}) = \mu$$
 $Var(\bar{X}) = \frac{\sigma^2}{n}$

In addition, the distribution of the sample mean is *normal* if *either* of the following is true:

- The underlying population is normal
- The sample size is large, say $n \ge 100$

Given the variance of the sample mean as $\sigma_{\bar{X}}^2 = \frac{\sigma^2}{n}$, its standard deviation, commonly referred to as the *standard error*, is $\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$.

Question 3: If the average of hourly wages in the population is $\mu = \$30$ and the variance of hourly wages is $\sigma^2 = 16$. Then what is the expected value, variance, standard error, and distribution of the sample mean estimator for a sample size of 400?

Question 4: You are interested in the average starting salary of CSUF graduates and are considering taking a random sample of 120 students. I advise you to take as large of a sample as feasible. This is sound advice because taking an even larger sample would ensure that

- \Box The sample average $\bar{x} = \mu$
- \Box The sample average \bar{x} is drawn from a normal distribution
- \Box The sample average \bar{x} is drawn from a distribution with lower variance

Note: I am using \bar{x} to denote a realization of \bar{X} .

Question 5: Can you explain intuitively why the variance of the sample mean increases with σ^2 and decreases with n?

Confidence Intervals

Let $z_{\alpha/2}$ be the *z*-value that leaves area $\alpha/2$ in the upper tail of the normal distribution. Then $1 - \alpha$ confidence interval is given by

$$\bar{x} \pm \underbrace{z_{\alpha/2} \frac{\sigma}{\sqrt{n}}}_{\text{Margin of Error}}$$

Question 6: Continuing with Question 3, say you took a sample of 400 individuals and found the average hourly wages in your sample of $\bar{x} = 26$. Create a 95% confidence interval for the true population mean.